

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 November 2002 (14.11.2002)

PCT

(10) International Publication Number  
**WO 02/089777 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 9/70**,  
31/381

(21) International Application Number: PCT/EP02/04975

(22) International Filing Date: 6 May 2002 (06.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
01111110.1 8 May 2001 (08.05.2001) EP

(71) Applicants: **SCHWARZ PHARMA AG** [DE/DE];  
Alfred-Nobel-Strasse 10, 40789 Monheim (DE). **LTS  
LOHMANN THERAPIE-SYSTEME AG** [DE/DE];  
Lohmannstrasse 2, 56626 Andernach (DE).

(72) Inventors: **LAUTERBACH, Thomas**; Am Rittersberg  
44, 40595 Düsseldorf (DE). **SCHACHT, Dietrich, Wil-  
helm**; Brahmstrasse 2, 50935 Köln (DE). **WOLFF,  
Hans-Michael**; Richard-Wagner-Strasse 2, 40789 Mon-  
heim (DE). **MÜLLER, Walter**; Hindenburgwall 12,  
56626 Andernach (DE).

(74) Agents: **HOFFMANN EITLE** et al.; Arabellastrasse 4,  
81925 München (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

**Published:**

- with international search report
- entirely in electronic form (except for this front page) and  
available upon request from the International Bureau

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: IMPROVED TRANSDERMAL THERAPEUTIC SYSTEM FOR THE TREATMENT OF PARKINSON'S DISEASE

(57) Abstract: This invention provides the use of a silicone-based transdermal therapeutic system having an area of 10 to 40 cm<sup>2</sup> and containing 0.1 to 3.15 mg / cm<sup>2</sup> of Rotigotine as active ingredient, for the preparation of an anti-Parkinson medicament which effects an improvement, compared to a placebo treatment, of the condition of human Parkinson patients, measured according to the Unified Parkinson's Disease Rating Scale (UPDRS) parts II and III, of 2 units or more following administration for a time period of at least 7 weeks.



**WO 02/089777 A1**

## IMPROVED TRANSDERMAL THERAPEUTIC SYSTEM FOR THE TREATMENT OF PARKINSON'S DISEASE

Description5 Field of the Invention

The present invention relates to an effective method for treating or alleviating symptoms of Parkinson's Disease, and the use of a Transdermal Therapeutic System (TTS) for  
10 delivering the dopamine receptor agonist rotigotine in a sufficient amount and at a sufficient rate to provide therapeutically effective treatment or alleviation of symptoms of Parkinson's disease.

15 Technical Background

Parkinson's disease is believed to be primarily caused by the degeneration of dopaminergic neurons in the substantia nigra. This, in effect, results in loss of tonic dopamine secretion  
20 and dopamine-related modulation of neuronal activity in the caudate nucleus, and thus in a deficiency of dopamine in certain brain regions. The resulting imbalance of the neurotransmitters acetylcholine and dopamine eventually results in disease-related symptoms. Although usually  
25 regarded as a motor system disorder, Parkinson's Disease is now considered to be a more complex disorder that involves both motor and nonmotor systems. This debilitating disease is characterized by major clinical features including tremor, bradykinesia, rigidity, dyskinesia, gait disturbances, and  
30 speech disorders. In some patients, dementia may accompany these symptoms. Involvement of the autonomic nerve system may produce orthostatic hypotension, paroxysmal flushing, problems with thermal regulation, constipation, and loss of bladder and sphincter control. Psychological disorders such  
35 as loss of motivation and depression may also accompany Parkinson's Disease.

Parkinson's Disease is primarily a disease of middle age and beyond, and it affects both men and women equally. The highest rate of occurrence of Parkinson's Disease is in the age group over 70 years old, where Parkinson's Disease exists in 1.5 to 2.5% of that population. The mean age at onset is between 58 and 62 years of age, and most patients develop Parkinson's Disease between the ages of 50 and 79. There are approximately 800,000 people in the United States alone with Parkinson's Disease.

Early motor deficits of Parkinson's Disease can be traced to incipient degeneration of nigral dopamine-releasing cells. This neuronal degeneration produces a defect in the dopaminergic pathway that connects the substantia nigra to the striatum. As the disease progresses, refractory motor, autonomic, and mental abnormalities may develop, which implies that there is progressive degeneration of striatal receptor mechanisms.

The clinical diagnosis of Parkinson's Disease is based on the presence of characteristic physical signs. The disease is known to be gradual in onset, slowly progressive, and variable in clinical manifestation. Evidence suggests that the striatal dopamine content declines to 20% below levels found in age-matched controls before symptoms occur.

Treatment of Parkinson's disease has been attempted with, inter alia, L-dopa (levodopa), which still is the gold standard for the therapy of Parkinson's Disease. Levodopa passes the blood-brain barrier as a precursor for dopamine and is then converted into dopamine in the brain. L-dopa improves the symptoms of Parkinson's Disease but may cause severe side effects. Moreover, the drug tends to lose its effectiveness after the first two to three years of


treatment. After five to six years, only 25% to 50% of patients maintain improvement.

Furthermore a major drawback of currently utilized therapies for Parkinson's Disease is the eventual manifestation of the "fluctuation syndrome", resulting in "all-or-none" conditions characterized by alternating "on" periods of mobility with dyskinesias and "off" periods with hypokinesia or akinesia. Patients who display unpredictable or erratic "on-off" phenomena with oral anti-Parkinson therapy have a predictable beneficial response to i.v. administration of L-dopa and other dopamine agonists, suggesting that fluctuations in plasma concentrations of drug are responsible for the "on-off" phenomena. The frequency of "on-off" fluctuations has also been improved by continuous infusions of the dopamine receptor agonists apomorphine and lisuride. However, this mode of administration is inconvenient. Therefore, other modes of administration providing a more constant plasma level, such as topical administration, are beneficial and have been suggested in the past.

As mentioned above, one treatment approach for Parkinson's disease involves dopamine receptor agonists. Dopamine receptor agonists (sometimes also referred to as dopamine agonists) are substances which, while structurally different from dopamine, bind to different subtypes of dopamine receptors and trigger an effect which is comparable to that of dopamine. Due to the reduced side-effects, it is advantageous when the substances selectively bind to a subgroup of dopamine receptors, i.e. the D2 receptors.

One dopamine receptor agonist which has been used to treat the symptoms of Parkinson's Disease is rotigotine. It has mostly been used in the form of its hydrochloride. Rotigotine is the International Non-Proprietary Name (INN) of the compound (-)-5,6,7,8-tetrahydro-6-[propyl]-[2-(2-

thienyl) ethyl] -amino] 1-naphthalenol having the structure shown below



5

To date, various transdermal therapeutic systems (TTS) for the administration of rotigotine have been described. WO 94/07468 discloses a transdermal therapeutic system containing rotigotine hydrochloride as active substance in a two-phase matrix which is essentially formed by a hydrophobic polymer material as the continuous phase and a disperse hydrophilic phase contained therein and mainly containing the drug and hydrated silica. The silica enhances the maximum possible loading of the TTS with the hydrophilic salt. Moreover, the formulation of WO 94/07468 usually contains additional hydrophobic solvents, permeation-promoting substances, dispersing agents and, in particular, an emulsifier which is required to emulsify the aqueous solution of the active principle in the lipophilic polymer phase. A TTS, prepared by using such a system, has been tested in healthy subjects and Parkinson patients. The average drug plasma levels obtained by using this system were around 0.15 ng/ml with a 20 cm<sup>2</sup> patch containing 10 mg rotigotine. This level must be considered as too low to achieve a truly efficacious treatment or alleviation of symptoms related to Parkinson's disease.

Various further transdermal therapeutic systems have been described in WO 99/49852. The TTS used in this patent

application comprise a backing layer, inert with respect to the constituents of the matrix, a self-adhesive matrix layer containing an effective quantity of rotigotine or rotigotine hydrochloride and a protective film which is to be removed before use. The matrix system is composed of a non-aqueous polymer adhesive system, based on acrylate or silicone, with a solubility of rotigotine of at least 5% w/w. Said matrix is essentially free of inorganic silicate particles. In Examples 1 and 2 and in Figure 1 of WO 99/49852 two transdermal therapeutic systems are compared. These are based on acrylate or silicone adhesives, respectively. Figure 1 of WO 99/49852 shows that a silicone patch releases about the same amount of active principle through skin as an acrylate patch. This has been demonstrated by the almost identical drug flux rates in an in vitro model, independent of the adhesive test system employed. Therefore an identical flux rate through human skin was expected.

It should be noted that the drug content of the silicone patch used in WO 99/49852 was lower than the drug content of the acrylate patch. However, this merely reflects the difference in solubility of the drug in the respective polymeric silicone and acrylate adhesives used in Examples 1 and 2, respectively. The TTS used in both examples contained the drug at about its saturation solubility in the respective adhesive systems. While the acrylate system is able to dissolve more drug than the silicone system, silicone in turn allows for a better release of the drug to skin. As these two effects compensate each other, it has been thought that the acrylate and the silicone systems as used in WO 99/49852 are about equivalent in the obtainable drug plasma levels and, hence, in therapeutic efficacy.

Considering the rather discouraging experiences made with the silicone formulation of WO 94/07568, the acrylate-based TTS of Example 1 of WO 99/49852 has been subjected to clinical

tests (safety and pharmacokinetic studies). The mean steady flux rate across human skin in vitro of this TTS amounted to 15.3  $\mu\text{g}/\text{cm}^2/\text{h}$ . However, it turned out that the plasma levels obtained using this TTS were still unsatisfactory and too low to allow for a really efficacious treatment of Parkinson's Disease. A 30 mg (20  $\text{cm}^2$ ) patch only yielded a mean maximum plasma concentration of 0.12 ng/ml, while a 5  $\text{cm}^2$  patch containing 7.5 mg yielded a mean maximum plasma concentration of 0.068 ng/ml. Again, such values have to be considered as too low to provide a real therapeutic progress in the treatment of Parkinson's Disease. Thus, in summary, both the 20  $\text{cm}^2$  silicone patch of WO 94/07468 and the 20  $\text{cm}^2$  acrylate patch of WO 99/49852 failed to evoke sufficient drug plasma levels to provide a satisfactory therapeutic effectiveness.

In view of these experiences, it has been very surprising that a transdermal therapeutic system containing rotigotine in free base form in a silicone matrix could not only provide unexpectedly high plasma levels of rotigotine but also a significant therapeutic progress in the transdermal treatment of Parkinson's Disease. In particular, it has been unexpected that a transdermal therapeutic system having a size of as little as 10 or 20  $\text{cm}^2$  could provide for an effective treatment of Parkinson's Disease in a placebo-controlled clinical study, as indicated by an improvement in the Unified Parkinson's Disease Rating Scale (UPDRS) of 2 or more compared to a placebo treatment. In the context of this application, "placebo treatment" refers to a treatment with a transdermal therapeutic system of identical qualitative composition and according to the same therapy regimen but where the active ingredient (rotigotine) in the transdermal therapeutic system has been omitted.

It should be understood that the term "treatment" in the context of this application is meant to designate a treatment or alleviation of the symptoms of Parkinson's Disease, rather

than a real causative treatment of Parkinson's Disease leading to a complete cure.

## 5 Summary of the Invention

The present invention provides the use of a silicone-based transdermal therapeutic system having an area of 10 to 40 cm<sup>2</sup> and containing 0.1 to 3.15 mg / cm<sup>2</sup> of Rotigotine as active  
10 ingredient, for the preparation of an anti-Parkinson medicament which effects an improvement, compared to a placebo treatment, of the condition of human Parkinson patients, measured according to the Unified Parkinson's Disease Rating Scale (UPDRS) parts II and III, of 2 units or  
15 more following administration for a time period of at least 7 weeks, preferably at least 11 weeks.

The silicone-based transdermal therapeutic system as used in the present invention must contain at least one amine  
20 resistant silicone compound as the main component. Usually, the silicone compound will be a pressure sensitive adhesive or a mixture thereof and will form a matrix in which the other components of the TTS are embedded. Moreover, the adhesive(s) should preferably be pharmaceutically acceptable  
25 in a sense that it is biocompatible, non-sensitizing and non-irritating to skin. Particularly advantageous silicone adhesives for use in the present invention should further meet the following requirements:

- Retained adhesive and cohesive properties in the presence  
30 of moisture or perspiration, under normal temperature variations,
- good compatibility with rotigotine as well as with the further excipients used in the formulation; in particular, the adhesive should not react with the amino group contained  
35 in rotigotine.



It has been shown that pressure sensitive adhesives of the type forming a soluble polycondensed polydimethylsiloxane (PDMS) / resin network, wherein the hydroxy endgroups are capped with e.g. trimethylsilyl (TMS) groups, are particularly useful in the practice of the present invention. Preferred adhesives of this kind are the BIO-PSA silicone pressure sensitive adhesives manufactured by Dow Corning, particularly the Q7-4201 and Q7-4301 qualities. However, other silicone adhesives may likewise be used.

In further and preferred aspects the present invention also provides a silicone-based transdermal therapeutic system comprising two or more silicone adhesives as the main adhesive components for such use. It can be advantageous if such a mixture of silicone adhesives comprises at least one high tack and at least one medium tack adhesive to provide for the optimum balance between good adhesion and little cold flux. Excessive cold flux may result in a too soft patch which easily adheres to the package or to patient garments. Moreover, such a mixture of adhesives seems to be particularly useful for obtaining an efficacious transdermal therapeutic system. A mixture of the aforementioned Q7-4201 (medium tack) and Q7-4301 (high tack) amine resistant silicone pressure sensitive adhesives in about equal amounts proved to be particularly useful in the practice of this invention.

In a further preferred embodiment, the silicone-based transdermal therapeutic system further includes a solubilizer. Several surfactant-like or amphiphilic substances may be used as solubilizers. They should be pharmaceutically acceptable and approved for use in medicaments. It is advantageous if the solubilizer also acts to improve the cohesion of the transdermal therapeutic system. A particularly preferred example of such a solubilizer is soluble polyvinylpyrrolidone.

Polyvinylpyrrolidone is commercially available, e.g. under the trademark Kollidon<sup>®</sup> (Bayer AG). Other examples include copolymers of polyvinylpyrrolidone and vinyl acetate, polyethyleneglycol, polypropylene glycol, glycerol and fatty acid esters of glycerol or copolymers of ethylene and vinylacetate.

The silicone-based transdermal therapeutic system for use according to the present invention preferably contains less than 1 wt% of inorganic silicates, most preferably it is completely free from inorganic silicates.

The water content in the transdermal therapeutic systems for use in the present invention is preferably low enough so that no evaporation of water during preparation of the TTS is necessary. Typically, the water content in a freshly prepared patch is below 2%, more preferably 1 wt% or lower.

In a particularly preferred embodiment of the present invention, the transdermal therapeutic system has a surface area of 10 to 30 cm<sup>2</sup>, more preferably 20 to 30 cm<sup>2</sup>. It goes without saying that a TTS having a surface area of, say, 20 cm<sup>2</sup> is pharmacologically equivalent to and may be exchanged by two 10 cm<sup>2</sup> patches or four 5 cm<sup>2</sup> patches having the same drug content per cm<sup>2</sup>. Thus, the surface areas as indicated in this application should be understood to refer to the total surface of all TTSSs simultaneously administered to a patient.

Providing and applying one or several transdermal therapeutic systems according to the invention has the pharmacological advantage over oral therapy that the attending physician can titrate the optimum dose for the individual patient relatively quickly and accurately, e.g. by simply increasing the number or size of patches given to the patient. Thus, the optimum individual dosage can often be determined after a time period of only about 3 weeks with low side effects.

A preferred content of rotigotine per patch is in the range of 0.1 to 2.0 mg / cm<sup>2</sup>. Still more preferred are 0.4 to 1.5 mg rotigotine per cm<sup>2</sup>. If a 7 day patch is desired, higher  
5 drug contents will generally be required. A rotigotine content in the range of about 0.4 to 0.5 wt% has been found to be particularly advantageous in that it provides the optimum usage of the drug contained in the TTS, i.e. there is only very little residual drug content in the TTS after  
10 administration. The apparent dose administered by using such a TTS usually is 50% or more and may be as high as 80-90% of the drug amount originally contained in the TTS.

The fact that the silicone-based transdermal therapeutic  
15 system described in this invention is able to provide a significant therapeutic effect against symptoms of Parkinson's Disease even at surface areas of 10 to 30 cm<sup>2</sup> and particularly as little as 10 or 20 cm<sup>2</sup> and at low drug contents of about 0.4 to 0.5 mg/cm<sup>2</sup>, particularly about 0.45  
20 g/cm<sup>2</sup>, must be considered as a further particular benefit provided by the present invention.

The transdermal therapeutic system used in the present invention usually is a patch having a continuous adhesive  
25 matrix in at least its center portion containing the drug. However, transdermal equivalents to such patches are likewise comprised by the present invention, e.g. an embodiment where the drug is in an inert but non-adhesive silicone matrix in the center portion of the TTS and is surrounded by an  
30 adhesive portion along the patch edges.

In a further aspect, this invention relates to a method of treating Parkinson's Disease by applying on a patient in need thereof a silicone-based transdermal therapeutic system  
35 having an area of 10 to 40 cm<sup>2</sup> and containing 0.1 to 3.15 mg / cm<sup>2</sup> of rotigotine as active ingredient, the improvement

wherein is that the condition of the patient, measured according to the Unified Parkinson's Disease Rating Scale (UPDRS) parts II and III, is improved, vis á vis placebo treatment, by about 2 units or more over a time period of at least 7 weeks administration. Maintenance of this UPDRS score improvement for up to 7 weeks has been shown as well. Thus, improving the UPDRS (parts II+III) score over placebo by at least 2 units following administration for 7, preferably 11 weeks forms a particularly beneficial aspect of the present invention.

Unless expressly indicated otherwise, any references to rotigotine in the context of this invention and the claims of this application mean rotigotine in the form of its free base. In some cases traces of rotigotine hydrochloride may be contained in rotigotine but these traces do typically not exceed 5 wt%, based on the amount of the free base. More preferably the content of hydrochloride impurities should be less than 2 wt%, even more preferably less than 1%, and most preferably the rotigotine used in the present invention contains less than 0.1 wt% or no hydrochloride impurities at all.

Extensive clinical trials using the transdermal therapeutic system described herein have shown that it is surprisingly possible to achieve and warrant a constant stimulation of the dopamine receptors of Parkinson patients, resulting in a notable improvement in the clinically relevant UPDRS for a time period of at least 7 weeks. More specifically, clinical studies using a patch according to the preparation example as given herein have resulted in the following improvements in the FAS UPDRS (part II and III) score following administration for 11 weeks:

Patch Size	Amount of rotigotine	Improvement in UPDRS over Placebo Treatment	p (one sided)
10 cm <sup>2</sup>	4.5 mg	- 2.148	0.0393
20 cm <sup>2</sup>	9.0 mg	- 3.123	0.0063
30 cm <sup>2</sup>	13.5 mg	- 4.909	0.0000
40 cm <sup>2</sup>	18.0 mg	- 5.035	0.0000

The abbreviation FAS stands for "Full Analysis Set" and thus designates an analysis including all patients who were included in the study. The UPDRS score and the study design is explained in more detail in the Clinical Trials Example below. Fig. 1 is a graph illustrating the mean change from base line in UPDRS (II + III) total scores from day 0 to end of treatment in this study. This figure compares the effects of a treatment according to the present invention with a placebo treatment. Statistically significant improvements can particularly be observed for patches having an area of 20 cm<sup>2</sup> or more, though even the effect of the 10 cm<sup>2</sup> patch must be considered as an improvement. The value indicated as p in the above table represents the one-sided p-value obtained from a statistical evaluation of the trial data.

While an improvement in the UPDRS scale of 2, compared to placebo, may already be called a success, an improvement of 3, 4 or even 5 or more units would even more represent a therapeutic progress and, as such, form a preferred aspect of the present invention.

Additional tests including pharmacokinetics, dose-activity relationship, compliance, and drug safety confirmed the therapeutic usefulness of the silicone-based transdermal therapeutic system used herein.

The invention and the best mode for carrying it out will be explained in more detail in the following non-limiting examples.

## 5 Preparation Example

A transdermal therapeutic system using a combination of silicone-type pressure sensitive adhesives was prepared as follows.

10

(-)-5,6,7,8-tetrahydro-6-[propyl-[2-(2-thienyl)ethyl]-amino]-1-naphthalenol hydrochloride (rotigotine hydrochloride, 150 g) was added to a solution of 17.05 g NaOH in 218 g ethanol (96%). The resulting mixture was stirred for approximately 15 10 minutes. Then 23.7 g of sodium phosphate buffer solution (8.35 g  $\text{Na}_2\text{HPO}_4 \times 2\text{H}_2\text{O}$  and 16.07 g  $\text{NaH}_2\text{PO}_4 \times 2\text{H}_2\text{O}$  in 90.3 g water) was added. Insoluble or precipitated solids were separated from the mixture by filtration. The filter was rinsed with 60.4 g ethanol (96%) to obtain a particle-free 20 ethanolic solution of rotigotine in the form of the free base.

The rotigotine free base solution (346.4 g) in ethanol (35% w/w) was mixed with 36.2 g ethanol (96%). The resulting 25 solution was mixed with 109 g of an ethanolic solution containing 25 wt% polyvinylpyrrolidone (KOLLIDON<sup>®</sup> 90F), 0.077 wt% aqueous sodium bisulfite solution (10 wt%), 0.25 wt% ascorbyl palmitate, and 0.63 wt% DL-alpha-tocopherol until homogenous. To the mixture, 817.2 g of an amine resistant 30 high tack silicone adhesive (BIO-PSA<sup>®</sup> Q7-4301 mfd. by Dow Corning) (74 wt% solution in heptane), 851.8 g of an amine resistant medium tack silicone adhesive (BIO-PSA<sup>®</sup> Q7-4201 mfd. by Dow Corning) (71 wt% solution in heptane), and 205.8 g petrol ether (heptane) were added, and all components were 35 stirred until a homogenous dispersion was obtained.

The dispersion was coated onto a suitable polyester release liner (SCOTCHPAK<sup>®</sup> 1022) with a suitable doctor knife and the solvents were continuously removed in a drying oven at temperatures up to 80°C for about 30 min to obtain a drug-containing adhesive matrix of 50 g/m<sup>2</sup> coating weight. The dried matrix film was laminated with a polyester-type backing foil (SCOTCHPAK<sup>®</sup> 1109). The individual patches were punched out of the complete laminate in the desired sizes (e.g. 10 cm<sup>2</sup>, 20 cm<sup>2</sup>, 30 cm<sup>2</sup>) and sealed into pouches under the flow of nitrogen.

The following table shows the composition in mg/20 cm<sup>2</sup> of a transdermal therapeutic system according to the present invention containing a combination of two silicone-type PSA.

15

Composition Components	Amount (mg)
Rotigotine Base	9.00
Polyvinylpyrrolidone	2.00
Silicone BIO-PSA <sup>®</sup> Q7-4301	44.47
Silicone BIO-PSA <sup>®</sup> Q7-4201	44.46
Ascorbyl palmitate	0.02
DL-alpha Tocopherol	0.05
Sodium metabisulfite	0.0006
Matrix coating weight	50 g/m <sup>2</sup>

### Clinical Trials

20 Methods:

The rotigotine TTS as prepared in the above preparation example has been tested in multicenter, placebo-controlled, double-blind, randomised clinical trials involving more than 300 Parkinson patients, and proven to yield an effective alleviation of the symptoms of Parkinson's Disease in patients suffering from this disease upon prolonged

administration times (11 weeks, of which 4 weeks were a titration period and 7 weeks were the maintenance period), once daily administration). The patients were not receiving other dopaminergic medications at that time.

5

After a 4-7 day open-label placebo run-in period, 329 patients were randomized to placebo or one of four rotigotine daily dosages (drug content in patch 4.5 mg, 9.0 mg, 13.5 mg or 18 mg), and followed for a 4-week dose-titration period, a 10 7-week dose-maintenance period, a 1-week dose de-escalation period and a 2-week safety follow-up period in a double-blind fashion. The study design (therapy plan) is illustrated in more detail in Fig. 2.

15 The prespecified primary efficacy outcome was a change in the Activities of Daily Living and Motor components of the generally accepted Unified Parkinson's Disease Rating Scale (UPDRS II/III) between baseline and the last evaluation on treatment (week 11).

20

Parts II and III of the UPDRS measure and rate the following clinical parameters in Parkinson patients:

## 25 II. ACTIVITIES OF DAILY LIVING

25

### A. Speech

0= Normal.

1= Mildly affected. No difficulty being understood.

30 2= Moderately affected. Sometimes asked to repeat statements.

3= Severely affected. Frequently asked to repeat statements.

4= Unintelligible most of the time.

35



## B. Salivation

0= Normal.

1= Slight but definite excess of saliva in mouth; may have  
5 nighttime drooling.

2= Moderately excessive saliva; may have minimal drooling.

3= Marked excess of saliva with some drooling.

4= Marked drooling, requires constant tissue or  
handkerchief.

10

## C. Swallowing

0= Normal.

15 1= Rare choking.

2= Occasional choking.

3= Requires soft food.

4= Requires NG tube or gastronomy feeding.

20

## D. Handwriting

0= Normal.

1= Slightly slow or small.

25 2= Moderately slow or small; all words are legible.

3= Severely affected; not all words are legible.

4= The majority of words are not legible.

## 30 E. Cutting food and handling utensils

0= Normal.

1= Somewhat slow and clumsy, but no help needed.

2= Can cut most foods, although clumsy and slow, some help  
35 needed.

3= Food must be cut by someone, but can still feed slowly.

4= Needs to be fed.

F. Dressing

5

0= Normal.

1= Somewhat slow, but no help needed.

2= Occasional assistance with buttoning, getting arms in sleeves.

10 3= Considerable help required, but can do some things alone.

4= Helpless.

15 G. Hygiene

0= Normal.

1= Somewhat slow, but no help needed.

20 2= Needs help to shower or bathe; or very slow in hygienic care.

3= Requires assistance for washing, brushing teeth, combing hair, going to bathroom.

4= Foley catheter or other mechanical aids.

25

H. Turning in bed and adjusting bed clothes

0= Normal.

1= Somewhat slow and clumsy, but no help needed.

30 2= Can turn alone or adjust sheets, but with great difficulty.

3= Can initiate, but not turn or adjust sheets alone.

4= Helpless.

35

I. Falling (unrelated to freezing)

- 0= None.  
1= Rare falling.  
2= Occasionally falls, less than once per day.  
5 3= Falls an average of once daily.  
4= Falls more than once daily.

J. Freezing when walking

10

- 0= None.  
1= Rare freezing when walking; may have start hesitation.  
2= Occasional freezing when walking.  
3= Frequent freezing. Occasionally falls from freezing.  
15 4= Frequent falls from freezing.

K. Walking

20

- 0= Normal.  
1= Mild difficulty. May not swing arms or may tend to drag leg.  
2= Moderate difficulty, but requires little or no assistance.  
25 3= Severe disturbance of walking, requiring assistance.  
4= Cannot walk at all, even with assistance.

L. Tremor (Symptomatic complaint of tremor in any part of  
30 body.)

- 0= Absent.  
1= Slight and infrequently present.  
2= Moderate; bothersome to patient.  
35 3= Severe; interferes with many activities.  
4= Marked; interferes with most activities.

## M. Sensory complaints related to parkinsonism

- 5 0= None.  
1= Occasionally has numbness, tingling, or mild aching.  
2= Frequently has numbness, tingling, or aching; not distressing.  
3= Frequent painful sensations.  
10 4= Excruciating pain.

## III. MOTOR EXAMINATION

## 15 N. Speech

- 0= Normal.  
1= Slight loss of expression, diction and/or volume.  
2= Monotone, slurred but understandable; moderately  
20 impaired.  
3= Marked impairment, difficult to understand.  
4= Unintelligible.

## 25 O. Facial Expression

- 0= Normal.  
1= Minimal hypomimia, could be normal "Poker Face".  
2= Slight but definitely abnormal diminution of facial  
30 expression.  
3= Moderate hypomimia; lips parted some of the time.  
4= Masked or fixed facies with severe or complete loss of facial expression; lips parted  $\frac{1}{4}$  inch or more.

35

## P. Tremor at rest (head, upper and lower extremities)

- 0= Absent.
- 1= Slight and infrequently present.
- 2= Mild in amplitude and persistent. Or moderate in
- 5 amplitude, but only intermittently present.
- 3= Moderate in amplitude and present most of the time.
- 4= Marked in amplitude and present most of the time.

10 Q. Action or Postural Tremor of hands

- 0= Absent.
- 1= Slight; present with action.
- 2= Moderate in amplitude; present with action.
- 15 3= Moderate in amplitude with posture holding as well as action.
- 4= Marked in amplitude; interferes with feeding.

20 R. Rigidity (Judged on passive movement of major joints with patient relaxed in sitting position. Cogwheeling to be ignored).

- 0= Absent.
- 25 1= Slight or detectable only when activated by mirror or other movements.
- 2= Mild to moderate.
- 3= Marked, but full range of motion easily achieved.
- 4= Severe, range of motion achieved with difficulty.

30

S. Finger Taps (Patient taps thumb with index finger in rapid succession.)

- 35 0= Normal.
- 1= Mild slowing and/or reduction in amplitude.

- 2= Moderately impaired. Definite and early fatiguing. May have occasional arrests in movement.
- 3= Severely impaired. Frequent hesitation in initiating movements or arrests in ongoing movement.
- 5 4= Can barely perform the task.
- T. Hand Movements (Patient opens and closes hands in rapid succession.)
- 10 0= Normal.
- 1= Mild slowing and/or reduction in amplitude.
- 2= Moderately impaired. Definite and early fatiguing. May have occasional arrests in movement.
- 3= Severely impaired. Frequent hesitation in initiating movements or arrests in ongoing movement.
- 15 4= Can barely perform the task.
- U. Rapid Alternating Movements of Hands (Pronation-supination movements of hands, vertically and horizontally, with as large an amplitude as possible, both hands simultaneously.)
- 20 0= Normal.
- 25 1= Mild slowing and/or reduction in amplitude.
- 2= Moderately impaired. Definite and early fatiguing. May have occasional arrests in movement.
- 3= Severely impaired. Frequent hesitation in initiating movements or arrests in ongoing movement.
- 30 4= Can barely perform the task.
- V. Leg Ability (Patient taps heel on the ground in rapid succession picking up entire leg. Amplitude should be at least 3 inches.)
- 35

- 0= Normal.
- 1= Mild slowing and/or reduction in amplitude.
- 2= Moderately impaired. Definite and early fatiguing. May have occasional arrests in movement.
- 5 3= Severely impaired. Frequent hesitation in initiating movements or arrests in ongoing movement.
- 4= Can barely perform the task.
- 
- 10 W. Arising from Chair (Patient attempts to rise from a straightbacked chair, with arms folded across chest.)
- 0= Normal.
- 1= Slow; or may need more than one attempt.
- 15 2= Pushes self up from arms of seat.
- 3= Tends to fall back and may have to try more than one time, but can get up without help.
- 4= Unable to arise without help.
- 
- 20 X. Posture
- 0= Normal erect.
- 1= Not quite erect, slightly stooped posture; could be normal for older person.
- 25 2= Moderately stooped posture, definitely abnormal; can be slightly leaning to one side.
- 3= Severely stooped posture with kyphosis; can be moderately leaning to one side.
- 30 4= Marked flexion with extreme abnormality of posture.
- 
- Y. Gait
- 35 0= Normal.

- 1= Walks slowly, may shuffle with short steps, but no festination (hastening steps) or propulsion.
- 2= Walks with difficulty, but requires little or no assistance; may have some festination, short steps, or propulsion.
- 5 3= Severe disturbance of gait, requiring assistance.
- 4= Cannot walk at all, even with assistance.
- Z. Postural Stability (Response to sudden, strong posterior displacement produced by pull on shoulders while patient erect with eyes open and feet slightly apart. Patient is prepared.)
- 10 0= Normal.
- 15 1= Retropulsion, but recovers unaided.
- 2= Absence of postural response; would fall if not caught by examiner.
- 3= Very unstable, tends to lose balance spontaneously.
- 4= Unable to stand without assistance.
- 20 AA. Body Bradykinesia and Hypokinesia (Combining slowness, hesitancy, decreased armswing, small amplitude, and poverty of movement in general.)
- 25 0= None.
- 1= Minimal slowness, giving movement a deliberate character; could be normal for some persons. Possibly reduced amplitude.
- 30 2= Mild degree of slowness and poverty of movement which is definitely abnormal. Alternatively, some reduced amplitude.
- 3= Moderate slowness, poverty or small amplitude of movement.
- 35 4= Marked slowness, poverty or small amplitude of movement.



The total UPDRS score is determined from the individual scores as follows:

First, a baseline value is determined for each patient participating in the study. This is done by summing up the individual scores of the UPDRS part II and III parameters at day 0, i.e. before treatment. Any determinations of the UPDRS score in the course of treatment will then be compared to this baseline value and changes relative to baseline value are recorded. Finally the mean improvement in UPDRS II+III at Day 77 (week 11) relative to baseline will be determined by averaging over all trial subjects. The resulting value is designated as the FAS (Full Analysis Set) Randomized Mean Change from Baseline Value in Total UPDRS Score (II + III) and is plotted as the y-axis in Figure 1. The term "randomized" refers to the fact that the patients were randomized to the different pre-defined doses in advance.

Patients suffering from Parkinson's Disease are known to experience a relatively strong placebo effect, i.e. even a placebo treatment improves the UPDRS score of Parkinson patients to some extent. It is therefore important to compare any effects of drug treatment with the level of UPDRS improvement attained by a placebo treatment of the same length. The final evaluation of improvement is therefore made relative to the effect of a placebo treatment of the same duration.

#### Results:

There was a significant, dose-related improvement in UPDRS II/III scores between baseline and week 11 when applying the TTS according to the present invention, particularly for the 9.0, 13.5, and 18 mg groups, compared to placebo. This result is apparent from Figure 1 and the following table:

Patch Size	Amount of rotigotine	Improvement in Mean Total UPDRS II+III over Placebo Treatment at week 11	p (one sided)
10 cm <sup>2</sup>	4.5 mg	- 2.148	0.0393
20 cm <sup>2</sup>	9.0 mg	- 3.123	0.0063
30 cm <sup>2</sup>	13.5 mg	- 4.909	0.0000
40 cm <sup>2</sup>	18.0 mg	- 5.035	0.0000

Rotigotine, when administered using the inventive TTS, was generally well tolerated. Application site skin reactions were generally mild and occurred with placebo patches, but were more common among subjects randomized to the higher dosage groups. There were no differences among the groups regarding vital signs, laboratory tests and ECGs.

#### Conclusions:

The above results show for the first time in a placebo-controlled study that a dopamine agonist (rotigotine), administered transdermally and once daily by a specific TTS, produces clinical improvement with satisfactory tolerability and safety in patients with early Parkinson's Disease.

Such a result could neither be obtained with the acrylic transdermal therapeutic system of WO 99/49852 nor with the silicone transdermal therapeutic system of WO 94/07468. Before this background, this result must be viewed as particularly surprising and beneficial for Parkinson patients.

## Claims

1. The use of a silicone-based transdermal therapeutic  
5 system having an area of 10 to 40 cm<sup>2</sup> and containing 0.1  
to 3.15 mg / cm<sup>2</sup> of rotigotine as active ingredient, for  
the preparation of an anti-Parkinson medicament which  
effects an improvement, vis á vis placebo treatment, in  
the condition of a human Parkinson patient, measured  
10 according to the Unified Parkinson's Disease Rating  
Scale (UPDRS) parts II and III, of 2 units or more  
following administration for at least 7 weeks.
2. The use according to claim 1 wherein the silicone-based  
15 transdermal therapeutic system comprises two or more  
silicone adhesives as the main adhesive components.
3. The use according to claim 1 or 2 wherein the silicone-  
based transdermal therapeutic system further includes a  
20 solubilizer.
4. The use according to claim 3 wherein the solubilizer is  
polyvinylpyrrolidone.
- 25 5. The use according to any of the preceding claims wherein  
the silicone-based transdermal therapeutic system  
contains less than 1 wt% of inorganic silicates.
6. The use according to claim 5 wherein the silicone-based  
30 transdermal therapeutic system is free from inorganic  
silicates.
7. The use according to any of the preceding claims wherein  
the transdermal therapeutic system has an area of 10 to  
35 30 cm<sup>2</sup>.

8. The use according to any of the preceding claims wherein the transdermal therapeutic system contains 0.1 to 1.5 mg / cm<sup>2</sup> of rotigotine.

5 9. The use according to claim 1 wherein the transdermal therapeutic system is a patch having an area of 10 to 30 cm<sup>2</sup> and a content of rotigotine of 0.4 to 0.5 mg/cm<sup>2</sup> in an adhesive silicone-based matrix.

10 10. In a method of treating Parkinson's Disease by applying on a patient suffering from this disease a silicone-based transdermal therapeutic system having an area of 10 to 40 cm<sup>2</sup> and containing 0.1 to 3.15 mg/cm<sup>2</sup> of rotigotine as active ingredient, the improvement wherein  
15 is that the condition of the patient, measured according to the Unified Parkinson's Disease Rating Scale (UPDRS) parts II and III, is improved, vis à vis placebo treatment, by about 2 units or more following administration for a time period of at least 7 weeks.

20

11. A transdermal therapeutic system for the treatment of Parkinson's Disease, which has a size of 10 to 30 cm<sup>2</sup> and contains 0.4 to 0.5 mg/cm<sup>2</sup> rotigotine as active ingredient in a matrix mainly comprising an admixture of  
25 at least two amine-resistant silicone adhesives.

12. The transdermal therapeutic system according to claim 11 which contains less than 1 wt% of inorganic silicates.

30 13. The transdermal therapeutic system according to claim 11 which further includes a solubilizer.

14. The transdermal therapeutic system of claim 13 wherein the solubilizer is polyvinylpyrrolidone.

35

1/2





Fig. 1

Fig. 2



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/04975

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K9/70 A61K31/381

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 49852 A (LOHMANN THERAPIE SYST LTS ;MUELLER WALTER (DE); PECK JAMES V (US);) 7 October 1999 (1999-10-07) cited in the application examples 2,6 page 7, paragraph 2 page 9, paragraph 2 -page 10, paragraph 1 -----	1-14
A	DE 199 40 238 A (LOHMANN THERAPIE SYST LTS) 1 March 2001 (2001-03-01) column 2, line 30 -column 2, line 48 column 3, line 32 -column 3, line 55 -----	1-14

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

8 July 2002

Date of mailing of the international search report

01/08/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Borst, M

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 02/04975

### Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —  
because they relate to subject matter not required to be searched by this Authority, namely:  

Although claim 10 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition (Rule 39.1(iv) PCT).
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/04975

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9949852	A	07-10-1999	DE 19814084 A1	14-10-1999
			AT 210973 T	15-01-2002
			AU 746856 B2	02-05-2002
			AU 2934199 A	18-10-1999
			BR 9909313 A	21-11-2000
			CA 2326630 A1	07-10-1999
			CN 1295470 T	16-05-2001
			DE 59900581 D1	31-01-2002
			DK 1033978 T3	15-04-2002
			WO 9949852 A1	07-10-1999
			EP 1033978 A1	13-09-2000
			HU 0101519 A2	28-09-2001
			JP 2002509878 T	02-04-2002
			NO 20004915 A	08-11-2000
			PL 343255 A1	30-07-2001
			SK 14462000 A3	09-04-2001
			TR 200002829 T2	22-01-2001
DE 19940238	A	01-03-2001	DE 19940238 A1	01-03-2001
			AU 6995700 A	19-03-2001
			WO 0113899 A2	01-03-2001
			EP 1143939 A2	17-10-2001